

Trauma Informed Care: Incorporating into Your Practice

Spring, 2025

Presenter:

Cori Green, MD, MS

Associate Professor, Clinical Pediatrics
Weill Cornell Medicine, New York-
Presbyterian Hospital

Disclosures

“Neither I nor my spouse/partner has a relevant financial relationship with a commercial interest to disclose.”

WHY is trauma informed care
important in primary care?

Trauma-Informed Care in Practice

- SAMHSA (2015) concept of a trauma-informed approach - A program, organization, or system that is trauma-informed:
 - *Realizes* the widespread impact of trauma and understands potential paths for recovery
 - *Recognizes* the signs and symptoms of trauma in clients, families, staff, and others involved with the system
 - *Responds* by fully integrating knowledge about trauma into policies, procedures, and practices
 - *Seeks* to actively resist *re-traumatization*

Trauma is COMMON
(estimates that more than
50% of children have had
exposure to a traumatic
event)

Many children/families who
have experienced trauma are
UNDIAGNOSED and
UNTREATED

Providers who use a trauma
informed approach in ALL
patient encounters are
providing a UNIVERSAL
PRECAUTION to prevent
retraumatizing a child/family

Rationale

Focus on:

- Recovery and healing are possible
 - neuroplasticity, neurogenesis
- Protective factors facilitate healing and resilience
- Healing takes place in the context of safe and supportive relationships

Why become trauma informed?

- Trauma is pervasive
- Impact is far-reaching
- Affects how people approach health care and other services
- Helping services can be inadvertently re-traumatizing

What are challenges incorporating trauma informed care into practice?

Change is a process

Incorporation into practice - challenges



Office culture (readiness for change)



Knowledge



Time/scheduling/workflow issues



Resources - educational, emotional, psychosocial, community



Ongoing staff support/team building



Financial barriers

Prevention and Promotion




Figure Legend:

A public health approach to prevent childhood toxic stress is a public health approach to promote relational health. Many of the components of a public health approach to prevent, mitigate, and treat toxic stress responses (see examples) are also components of a public health approach to promote, identify barriers to, and repair SSNRs. The examples provided are illustrative and not intended to be comprehensive or exhaustive. See the Appendix for full descriptions of the abbreviations. BStC, biological sensitivity to context; PTSD, posttraumatic stress disorder. Adapted with permission from Garner AS, Saul RA. Thinking Developmentally: Nurturing Wellness in Childhood to Promote Lifelong Health. Itasca, IL: American Academy of Pediatrics; 2018


7 C's of Resilience



COMPETENCE



CONFIDENCE



CONNECTION




CHARACTER



CONTRIBUTION







COPING



CONTROL

Positive Childhood Experiences Protective



Surveillance - trauma is common

“Has anything bad or scary or upsetting happened since our last visit?”



“Has anyone come or gone from your family recently?”

Initial Steps

Consider the diagnosis of trauma in patients with

Somatic complaints

Symptoms of
emotional
dysregulation/distress

Children living in
“high risk” situations

Sleeping/eating
difficulties

Academic/social
difficulties

Regression,
behavioral changes

Rule out co-morbidities (full assessment)

Utilize	Utilize Standardized Screening Tools
Assess	Assess degree of impairment
Address	Address safety issues

Screening

ACEs for parent and/or child

Domestic violence screening

SEEK (Safe Environment for Every Kid)

PCPs play a crucial role in trauma treatment

Childhood exposure to traumatic events is common

PCP offices can be the first place it is identified

PCP can learn to recognize and treat acute symptoms in their offices

Successful treatment is built upon a strong and caring relationship with at least one adult

Office based intervention - psychoeducation

Emphasize the role of a single nurturing relationship in recovery from trauma

Discuss the biological basis of fear and its manifestations (fight/flight)

Relationships, routine and regulation

Offer some practical tips on ways to integrate these concepts into family life now

Introduce role of therapy for ongoing trauma treatment, and provide linkage and warm hand-off (if possible)

TIC goals in primary care

Convey	Convey hope
Build	Build resilience
Engage	Engage learning/thinking brain
Develop	Develop child's self-efficacy
Teach	Teach skills of self-regulation

Case: Eli, 8 years old

CC: “angry and moody whenever I ask him to do anything”

- Comes with stepmother. He lived with birth mother until he was 7. Removed by CPS for neglect
- Initially eager to please and easygoing. Loving towards 2-year-old stepsister.
- Now gets angry and yells at his stepmother when she tells him to do things
- Some nights hears him crying softly in his room
- Above average grades and likes to read. One friend in his class.
- Stepfather works 12-hour days
- On exam: Well groomed, down cast eyes, sad affect, shame when stepmother speaks, only smiles when step sib brings him a toy.

Initial Observations

- Mom looks frustrated, overwhelmed, and angry
- Child - sad, ashamed, quiet
- What else do you want to know?

Table Exercise: Assessment and Differential Diagnosis

- What else do you want to know?
 - What would you ask the stepmother?
 - What would you ask Eli?
 - How would you phrase the questions?
- What is on your differential diagnosis?

Differential diagnosis?

- Depression
- Anxiety
- Trauma
- Adhd
- ODD
- Aggression

Next step?



Interview Eli

Interview with Eli

Safety

Calm voice,
eye contact,
warmth

Convey hope,
empathy,
“helper”

Gentle
questioning

Recognize,
compliment
strengths

Be alert for s/s
of distress

Interview with Eli

- Admits he has nightmares about being sent back to his birth mother
- Describes times when his birthmother or her friends hurt him.
- Admits he has thought of running away from home.
- CATS 2 score - 25

Initial Assessment: Eli

Child's trauma
screen
positive

Safety
concerns

Maternal
distress

Child - mood
and sleep
impairments

Impairment of
parent/child
relationship

Additional
screens
pending

Next step?



Psychoeducation: How to discuss trauma with mom and child

What is toxic stress and how does the body react

Message to parents



Distinguish between normal stress and toxic stress

Occurs when a child feels unsafe over a long time and lacks the buffering effect of being loved and cared for by a supportive adult

Children who experience this level of stress have difficulty with managing their emotions, regulating sleep and appetite and developing relationships

Treatment is available and effective. The strongest predictor of recovery is the development of a supportive, loving relationship with at least one caring adult.

Positive experiences mitigate adverse experiences

Talking with families about trauma

Describe	Describe signs and symptoms of childhood trauma (fight/flight/freeze response)
Convey	Convey support and hope for recovery, and assure collaboration and partnership
Offer	Offer practical suggestions for help NOW with difficult symptoms
Plant	Plant the seed of psychotherapy (begin linkage if family is ready)

- Might

**What might you recommend
for Eli's anger/mood issues?**

Self- regulation strategies

Parent helps child to label emotions, feelings, thoughts, and expand emotional vocabulary

Teach child about his brain - feeling, thoughts, behavior triangle

“Feelings thermometer”

Positive self-talk


Breathing exercises (box breathing)

Mindful meditation

FEELINGS SCALE

WHAT CAN YOU DO?





What office-based interventions would be helpful for Eli?

Sleep strategies

Self-regulation techniques

Relationship building/mom and child

Sleep interventions

Regular bedtime routine (bath, brush teeth, bedtime story, etc)

Relaxation techniques (gentle touch, box breathing, guided imagery), co-regulation

Mom sits with child in bedroom quietly, lighting low, attachment object

Mom checks-in at regular intervals after leaving

Sound machine?

Interventions for “strong emotions”

1

Reassurance - verbal and nonverbal messages to buffer child's fears

2

Routines - predictability, eg. school mornings, after school, bedtime

3

Regulation - Help child label emotions/ thoughts, “Time-in” activities, help with household tasks, play and co-regulation activities with parent

Resources

Aap.org: The Trauma-Informed Pediatric Practice. Paperback 2024

Aap.org: Childhood Trauma and Resilience : A practical guide. e-book. 2021

Thank you!